


Amendments to the Claims:

This listing of claims will replace all prior versions, and listing, of claims in the application:

Listing of Claims:

Claims 1-19. (Canceled)


Claim 20. (Previously presented) A method for treating phospholipase D (PLD) initiated polymorphoneutrophil (PMN) inflammation in a subject, comprising administering to the subject an effective anti-inflammatory amount of a lipoxin analog having the formula



wherein X is R₁, OR₁, or SR₁;

wherein R₁ is

- (i) a hydrogen atom;
- (ii) an alkyl of 1 to 8 carbons atoms, inclusive, which may be straight chain or branched;
- (iii) a cycloalkyl of 3 to 10 carbon atoms;
- (iv) an aralkyl of 7 to 12 carbon atoms;
- (v) phenyl;
- (vi) substituted phenyl




wherein Z_i , Z_{ii} , Z_{iii} , Z_{iv} and Z_v are each independently selected from $-\text{NO}_2$, $-\text{CN}$, $-\text{C}(=\text{O})-\text{R}_T$, $-\text{SO}_3\text{H}$, a hydrogen atom, halogen, methyl, $-\text{OR}_x$, wherein R_x is 1 to 8 carbon atoms, inclusive, which may be a straight chain or branched, and hydroxyl;

- (vii) a detectable label molecule; or
- (viii) a straight or branched chain alkenyl of 2 to 8 carbon atoms, inclusive;

wherein R_T is

- (i) a hydrogen atom;
- (ii) an alkyl of 1 to 8 carbons atoms, inclusive, which may be straight chain or branched;
- (iii) a cycloalkyl of 3 to 10 carbon atoms;
- (iv) an aralkyl of 7 to 12 carbon atoms;
- (v) phenyl;
- (vi) substituted phenyl



wherein Z_i , Z_{ii} , Z_{iii} , Z_{iv} and Z_v are each independently selected from $-\text{NO}_2$, $-\text{CN}$, $-\text{SO}_3\text{H}$, a hydrogen atom, halogen, methyl, $-\text{OR}_x$, wherein R_x is 1 to 8 carbon atoms, inclusive, which may be a straight chain or branched, and hydroxyl;

wherein Q_1 is $(\text{C}=\text{O})$, SO_2 or (CN) , provided when Q_1 is CN , then X is absent;


wherein one of R_2 and R_3 is a hydrogen atom and the other is

- (a) H;
- (b) an alkyl of 1 to 8 carbon atoms, inclusive, which may be a straight chain or branched;
- (c) a cycloalkyl of 3 to 6 carbon atoms, inclusive;
- (d) an alkenyl of 2 to 8 carbon atoms, inclusive, which may be straight chain or branched; or
- (e) $R_a Q_2 R_b$ wherein Q_2 is $-O-$ or $-S-$; wherein R_a is alkylene of 0 to 6 carbons atoms, inclusive, which may be straight chain or branched and wherein R_b is alkyl of 0 to 8 carbon atoms, inclusive, which may be straight chain or branched, provided when R_b is 0, then R_b is a hydrogen atom;

wherein R_4 is

- (a) H;
- (b) an alkyl of 1 to 6 carbon atoms, inclusive, which may be a straight chain or branched;

wherein R_5 is



wherein Z_i , Z_{ii} , Z_{iii} , Z_{iv} and Z_v are each independently selected from $-NO_2$, $-CN$, $-C(=O)-R_1$, $-SO_3H$, a hydrogen atom, halogen, methyl, $-OR_x$, wherein R_x is 1 to 8 carbon atoms, inclusive, which may be a straight chain or branched, and hydroxyl or a substituted or unsubstituted, branched or unbranched alkyl group;

wherein Y_1 is $-OH$, methyl, $-SH$, an alkyl of 2 to 4 carbon atoms, inclusive, straight chain or branched, an alkoxy of 1 to 4 carbon atoms, inclusive, or $CH_a Z_b$ where $a+b=3$, $a=0$ to 3, $b=0$ to 3 and Z is cyano, nitro or a halogen;

wherein R_6 is

- (a) H;

(b) an alkyl from 1 to 4 carbon atoms, inclusive, straight chain or branched;


wherein T is O or S, and pharmaceutically acceptable salts thereof, such that PLD initiated polymorphoneutrophil (PMN) inflammation is treated in a subject.

Claim 21. (Original): The method of claim 20, wherein said method is performed *in vitro*.

Claim 22. (Original): The method of claim 20, wherein said method is performed *in vivo*.

Claims 23-25. (Canceled)


Claim 26. (Previously presented) A method for treating phospholipase D (PLD) initiated superoxide generation or degranulation in a subject, comprising administering to the subject an effective anti-PLD amount of a lipoxin analog having the formula



wherein X is R₁, OR₁, or SR₁;

wherein R₁ is

- (i) a hydrogen atom;
- (ii) an alkyl of 1 to 8 carbons atoms, inclusive, which may be straight chain or branched;
- (iii) a cycloalkyl of 3 to 10 carbon atoms;
- (iv) an aralkyl of 7 to 12 carbon atoms;
- (v) phenyl;
- (vi) substituted phenyl




wherein Z_i , Z_{ii} , Z_{iii} , Z_{iv} and Z_v are each independently selected from $-\text{NO}_2$, $-\text{CN}$, $-\text{C}(=\text{O})-\text{R}_T$, $-\text{SO}_3\text{H}$, a hydrogen atom, halogen, methyl, $-\text{OR}_x$, wherein R_x is 1 to 8 carbon atoms, inclusive, which may be a straight chain or branched, and hydroxyl;

- (vii) a detectable label molecule; or
- (viii) a straight or branched chain alkenyl of 2 to 8 carbon atoms, inclusive;

wherein R_T is

- (i) a hydrogen atom;
- (ii) an alkyl of 1 to 8 carbons atoms, inclusive, which may be straight chain or branched;
- (iii) a cycloalkyl of 3 to 10 carbon atoms;
- (iv) an aralkyl of 7 to 12 carbon atoms;
- (v) phenyl;
- (vi) substituted phenyl



wherein Z_i , Z_{ii} , Z_{iii} , Z_{iv} and Z_v are each independently selected from $-\text{NO}_2$, $-\text{CN}$, $-\text{SO}_3\text{H}$, a hydrogen atom, halogen, methyl, $-\text{OR}_x$, wherein R_x is 1 to 8 carbon atoms, inclusive, which may be a straight chain or branched, and hydroxyl;

- (vii) a detectable label molecule; or
- (viii) a straight or branched chain alkenyl of 2 to 8 carbon atoms, inclusive;

wherein Q_1 is (C=O), SO_2 or (CN), provided when Q_1 is CN, then X is absent;


wherein one of R_2 and R_3 is a hydrogen atom and the other is

- (a) H;
- (b) an alkyl of 1 to 8 carbon atoms, inclusive, which may be a straight chain or branched;
- (c) a cycloalkyl of 3 to 6 carbon atoms, inclusive;
- (d) an alkenyl of 2 to 8 carbon atoms, inclusive, which may be straight chain or branched; or
- (e) $R_a Q_2 R_b$ wherein Q_2 is -O- or -S-; wherein R_a is alkylene of 0 to 6 carbons atoms, inclusive, which may be straight chain or branched and wherein R_b is alkyl of 0 to 8 carbon atoms, inclusive, which may be straight chain or branched, provided when R_b is 0, then R_b is a hydrogen atom;

wherein R_4 is

- (a) H;
- (b) an alkyl of 1 to 6 carbon atoms, inclusive, which may be a straight chain or branched;

wherein R_5 is



wherein Z_i , Z_{ii} , Z_{iii} , Z_{iv} and Z_v are each independently selected from $-NO_2$, $-CN$, $-C(=O)-R_1$, $-SO_3H$, a hydrogen atom, halogen, methyl, $-OR_x$, wherein R_x is 1 to 8 carbon atoms, inclusive, which may be a straight chain or branched, and hydroxyl or a substituted or unsubstituted, branched or unbranched alkyl group;

wherein Y_1 is -OH, methyl, -SH, an alkyl of 2 to 4 carbon atoms, inclusive, straight chain or

branched, an alkoxy of 1 to 4 carbon atoms, inclusive, or CH_aZ_b where $a+b=3$, $a=0$ to 3, $b=0$ to 3 and Z is cyano, nitro or a halogen;

wherein R_6 is

- (a) H;
- (b) an alkyl from 1 to 4 carbon atoms, inclusive, straight chain or branched;

wherein T is O or S, and pharmaceutically acceptable salts thereof, such that PLD initiated superoxide generation or granulation is treated in a subject.

Claim 27. (Original): The method of claim 26, wherein said method is performed *in vitro*.

Claim 28. (Original): The method of claim 26, wherein said method is performed *in vivo*.

Claim 29. (Canceled)

Claim 30. (Canceled)

Claim 31. (Canceled)

Claim 32. (Canceled)